

Exercises on Functions and Relations

Question 1: Consider the poset (A, \subseteq) , where $A = \{\{1\}, \{1, 2\}, \{1, 3\}, \{1, 2, 4\}\}$.

- (a) Find all maximum, minimum, maximal, and minimal elements of A .
- (b) Draw the Hasse Diagram of A .

Question 2:

- (a) Prove that the function $f : (\mathbb{N} \cup \{0\}) \times (\mathbb{N} \cup \{0\}) \longrightarrow \mathbb{N} \cup \{0\}$, defined by $f(k, n) = 2^k(2n + 1) - 1$ is one-to-one.
- (b) Prove that the function $f : (\mathbb{N} \cup \{0\}) \times (\mathbb{N} \cup \{0\}) \longrightarrow \mathbb{N} \cup \{0\}$, defined by $f(k, n) = 2^k(2n + 1) - 1$ is onto. (*You have to be very careful here.*)

Notice that parts (a) and (b) prove that \mathbb{N} has the same cardinality as $\mathbb{N} \times \mathbb{N}$, because the function f defined above is bijective (i.e. one-to-one and onto.)

- (c) Use (a) and (b) to prove that if A and B are countably infinite sets, then so is $A \times B$.

Question 3:

- (a) Prove that if A and B are both countably infinite disjoint sets, then so is $A \cup B$. *Hint: Consider the function $f : \mathbb{N} \longrightarrow A \cup B$, which maps even natural numbers to A and odd natural numbers to B . All you need to do is to give a mathematical formula for f .*

- (b) Prove that the intervals $(0,3)$ and $(3, \infty)$ have the same cardinality. *Hint: First show that $(0,3)$ and $(0,1)$ have the same cardinality by finding a one-to-one and onto function between them. Then show that $(3, \infty)$ has the same cardinality as \mathbb{R} by finding a one-to-one and onto function between them. Finally, depend on the fact that*

$(0,1)$ and \mathbb{R} have the same cardinality. When you come up with the two functions mentioned above, you'll have the following:

$(0,3)$ has the same cardinality as $(0,1)$, $(0,1)$ has the same cardinality as \mathbb{R} , \mathbb{R} has the same cardinality as $(3, \infty)$.

(c) Prove that the set of irrational numbers (i.e. $\mathbb{R} \setminus \mathbb{Q}$) is uncountable.

Notice that (c) implies that the cardinality of the set of irrational numbers is greater than the cardinality of the set of rational numbers.

Hints: Remember

1. If $f : A \rightarrow B$ is a bijective function, then $f^{-1} : B \rightarrow A$ is defined and it is also bijective.
2. If $f : A \rightarrow B$ and $g : B \rightarrow C$ are bijective functions, then $g \circ f : A \rightarrow C$ is also bijective.

Question 4:

- (1) Find a bijective function from $\mathbb{N} \times \mathbb{N}$ onto \mathbb{Z} .
- (2) Find a bijective function from $(-1, 1)$ onto $(\frac{9999}{777}, \infty)$.
- (3) Give a mathematical formula for a bijective function from \mathbb{Z} onto \mathbb{N} .
- (4) Find a bijective function from $\mathbb{N} \setminus \{1, 2, 3\}$ onto $(2\mathbb{N} - 1) \cup \{a, b, c\}$, where a , b , and c , are the letters a , b , c (This means $\mathbb{N} \cap \{a, b, c\} = \emptyset$.)